

INSTRUCTIONS



INHALATION NARCOSIS CHAMBER

Catalog No. 46420-0002



The Inhalation Narcosis Chamber is used to anesthetize small laboratory animals such as mice, hamsters, rats and guinea pigs.

1. Open the stopcock to allow entrance of fresh air into the chamber. This helps to prevent suffocation of the animal.
2. Raise the clear lid by releasing the Velcro closure.
3. Reach in and remove the perforated platform. Note that the platform folds in the center to allow it to fit through the top opening.
4. Place a pad of absorbent cotton about the size of the palm of your hand into the chamber bottom to absorb the anesthetic.
5. Replace the perforated platform by folding it, inserting it through the end and flattening it against the ledge.
6. Insert the animal into the chamber and close the lid with Velcro closure.
7. Pour anesthetic into the funnel sufficient for the size of animal in the chamber.
8. At the end of the work, remove the cotton from the chamber and dispose of it in an approved manner. Close the lid and the stopcock

Bel-Art Products

WAYNE, NJ 07470 USA

TEL: 1-800-4BEL-ART • FAX: 973-694-7199 • WWW.BELART.COM

946420013

Scienceware®, is a reg. TM of Bel-Art Products
Bel-Art Products assumes no obligation or liability for any advice furnished by it, or for results obtained with respect to these products. All such advice is given and accepted at the buyer's risk.

10/09

INSTRUCTIONS



INHALATION NARCOSIS CHAMBER

Catalog No. 46420-0002



The Inhalation Narcosis Chamber is used to anesthetize small laboratory animals such as mice, hamsters, rats and guinea pigs.

1. Open the stopcock to allow entrance of fresh air into the chamber. This helps to prevent suffocation of the animal.
2. Raise the clear lid by releasing the Velcro closure.
3. Reach in and remove the perforated platform. Note that the platform folds in the center to allow it to fit through the top opening.
4. Place a pad of absorbent cotton about the size of the palm of your hand into the chamber bottom to absorb the anesthetic.
5. Replace the perforated platform by folding it, inserting it through the end and flattening it against the ledge.
6. Insert the animal into the chamber and close the lid with Velcro closure.
7. Pour anesthetic into the funnel sufficient for the size of animal in the chamber.
8. At the end of the work, remove the cotton from the chamber and dispose of it in an approved manner. Close the lid and the stopcock

Bel-Art Products

WAYNE, NJ 07470 USA

TEL: 1-800-4BEL-ART • FAX: 973-694-7199 • WWW.BELART.COM

946420013

Scienceware®, is a reg. TM of Bel-Art Products
Bel-Art Products assumes no obligation or liability for any advice furnished by it, or for results obtained with respect to these products. All such advice is given and accepted at the buyer's risk.

10/09